

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

If swallowed this material may reduce the clotting ability of the blood and cause bleeding. Keep away from children, pets (especially dogs), and domestic animals. Do not contaminate feed and foodstuffs. Use only in rodent infested areas where exposure to agricultural crops or commodities will not occur. Wash thoroughly with soap and water after handling and before eating or smoking. Clean clothing should be used daily.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Bait exposed to soil surfaces may be hazardous to birds and other wildlife. Do not apply directly to lakes, streams, or ponds.

STORAGE AND DISPOSAL

Store in a locked enclosure (in a properly posted area) in such a manner so as not to present a hazard to persons, animals, food, feed, crops, or property. Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

5/26/93

**RODENT BAIT DIPHACINONE
TREATED GRAIN (0.01%)**

Active Ingredients: Diphacinone 2-diphenylacetyl-1, 3-indandione:	0.01%
Inert Ingredients:	99.99%
Total:	100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat. Do not induce vomiting or give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN

Intramuscular and oral administration of Vitamin K₁, combined with blood transfusions is indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin (Dicumarol).

REGISTRANT: California Department of Food and Agriculture
1220 "N" Street, Room A357
Sacramento, CA 95814

CALIF. REG. NO. 10965-50003
EPA SLN. NO. CA 890022
EPA EST. NO.

NET CONTENTS: _____

DIRECTIONS FOR USE

It is a violation of State and Federal laws to use this product inconsistent with its labeling.

**FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA
USE RESTRICTIONS:**

For use against GROUND SQUIRRELS, (*Spermophilus beecheyi*, *S. beldingi*), DEER MICE (*Peromyscus spp.*) and HOUSE MICE (*Mus musculus*).

IMPORTANT

Bait must be eaten at several feedings on five or more successive days with no period longer than 48 hours between feedings. Do not allow treated bait to contact skin. Use measuring spoon to fill bait boxes. Keep this product in a closed container, free from odors which may reduce acceptability.

BAITING: GROUND SQUIRRELS *Spermophilus beecheyi*, *S. beldingi*.

Scatter a handful of bait (about 10 baits per pound) evenly over 40 to 50 square feet near active burrows or runways. Retreat every other day for 3 to 4 applications. Do not pile bait!! An uninterrupted supply of bait should be available for 6 to 8 days.

BAITING: DEER MICE, *Peromyscus spp.* Spread bait evenly by hand, mechanical spreader, or aircraft at the rate of 2 to 6 pounds per acre through the infested area. Sufficient bait should be applied to last a minimum of 4 days (preferably without rain)

BAITING: HOUSE MICE, *Mus musculus*. Place tablespoon amounts (1/4 to 1/2 ounces) of bait in bait box or shallow container preferably in protected feeder stations at 8 to 12 foot intervals in dry locations such as in concealed places in corners or along walls where house mice feed, drink, or frequent. Inspect stations daily and add bait as needed. Replace moldy or old bait with fresh bait. Dispose of bait in such a manner as to prevent ingestion by children, pets, wildlife and domestic animals. An uninterrupted supply of bait should be maintained as long as any bait is taken which may be 2 to 4 weeks. Where continuous source of infestation is present permanent bait stations should be established and the bit replenished as needed. All poison baits must be picked up or disposed of after rodent activity has stopped.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Keep away from children, pets (especially dogs), and domestic animals. Do not contaminate feed and food stuffs. Use only in rodent-infested areas where exposure to agricultural crops or commodities will not occur. Wash thoroughly with soap and water after handling and before eating or smoking. Clean clothing should be used daily.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Bait exposed to soil surfaces may be hazardous to birds and other wildlife. Do not apply directly to lakes, streams, or ponds.

STORAGE AND DISPOSAL

Store in a locked enclosure (in a properly posted area) in such a manner so as not to present a hazard to persons, animals, food, feed, crops, or property. Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Directions for Use Continued

BAITING: WOOD RATS, *Neotoma* spp.

Place 4 to 6 ounces of bait in an open bait box or shallow container. Bait stations should be located near existing wood rat runways or dens, and generally spaced no further than 100 feet apart.

BAITING: COTTONTAIL RABBITS Orange County only.

Only for use in the production of an agricultural commodity or under Food and Agriculture Code Sections 6021-6024. Place 1 to 5 pounds of bait in a covered self-dispensing feeder or nursery flat in areas frequented by cottontail rabbits (near runways, resting or feeding areas). It may be necessary to move feeders to new locations to achieve bait acceptance. Carcasses of poisoned rabbits should be collected and disposed of in a sanitary landfill or by deep burying.

**RODENT BAIT DIPHACINONE
TREATED GRAIN (0.005%)**

Active Ingredients: Diphacinone 2-diphenylacetyl-1, 3-indandione:	0.005%
Inert Ingredients:	99.995%
Total:	100.000 %

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat. Do not induce vomiting or give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN

Intramuscular and oral administration of Vitamin K₁, combined with blood transfusions is indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin (Dicumarol).

REGISTRANT: California Department of Food and Agriculture
1220 "N" Street, Room A357
Sacramento, CA 94271-0001

CALIF. REG. NO. 10965-50001
EPA SLN. NO. CA 890020
EPA EST. NO.

NET CONTENTS: _____

DIRECTIONS FOR USE

It is a violation of State and Federal laws to use this product inconsistent with its labeling.

FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA

USE RESTRICTIONS:

For use against NORWAY RATS, ROOF RATS, GROUND SQUIRRELS, CHIPMUNKS, MUSKRATS, JACKRABBITS, MEADOW MICE, WOOD RATS, GOLDEN-MANTLED GROUND SQUIRRELS, and (COTTONTAIL RABBITS Orange County only).

IMPORTANT

An uninterrupted supply of bait should be maintained as long as any bait is taken, which may be 1 to 4 weeks. Treated baits should be placed in locations not accessible to children, pets, wildlife, and domestic animals, or in tamper-resistant bait boxes. Use a measuring scoop to fill bait boxes. Do not allow treated bait to contact skin. Initial acceptance may not occur until animals become accustomed to the bait station. Bait stations should have entrance holes large enough to admit target species, but not larger animals. Secure bait stations to prevent overturning.

Maintain bait supply for at least 15 days. Continue baiting until all signs of feeding have been stopped. Bait placements should be checked daily; replace consumed, contaminated, or spoiled baits.

Where a continuous source of infestation is present, permanent bait stations should be established and the bait replenished as needed. All poison baits must be picked up and disposed of after rodent activity has stopped. Keep this product in a closed container to maintain acceptability.

BAITING: NORWAY RATS, *Rattus norvegicus*, and ROOF RATS, *R. rattus*.

Place 4 to 16 ounces of bait in bait box or shallow container, preferable in protected feeder stations. Place bait stations in dry locations such as in concealed places, in corners, along walls where rats feed, drink or frequent. For roof rats, place bait at ground floor and top floor, or attic levels. For Norway rats, put bait at or near ground level and at burrows and harborages.

BAITING: GROUND SQUIRRELS, *Spermophilus beecheyi*, *S. beldingi*.

Bait boxes: Place 1 to 5 pounds of bait in a covered bait box in areas frequented by ground squirrels (near runways, burrows, etc.).

BAITING: GOLDEN-MANTLED GROUND SQUIRRELS, *Spermophilus lateralis*, and CHIPMUNKS, *Eutamias* spp.

Bait boxes: Place 1 to 5 pounds of bait in a covered bait box in areas frequented by target animals (near runways, burrows, etc.).

BAITING: JACK RABBITS (Black-tailed hare), *Lepus californicus*. Place 1 to 5 pounds of bait in a covered self-dispensing feeder or nursery flat in areas frequented by jack rabbits (near runways, resting or feeding areas). It may be necessary to move feeders to new locations to achieve bait acceptance. Carcasses of poisoned rabbits should be picked up and disposed of by deep burying or burning.

BAITING: MUSKRATS, *Ondatra zibethica*.

Floating bait boxes: Place 1 to 5 pounds of bait in floating bait box.

BAITING: MEADOW MICE, *Microtus californicus*, *Microtus montanus*.

Lightly scatter tablespoon amounts (1/4 to 1/2 ounce) of bait near active burrows or in runways. Repeat treatment every other day for three treatments.

FIFRA SECTION 2(ee) RECOMMENDATION FOR USE
IN THE STATE OF CALIFORNIA

CDFA RODENT BAIT BLOCK DIPHACINONE

EPA Reg. No. 61282-26-10965

Use Directions for Muskrats (*Ondatra zibethica*)

It is a violation of federal law to use this product in any manner inconsistent with its labeling. This recommendation is permitted under FIFRA Section 2(ee). This supplemental label and the federally registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on the EPA registered label.

DIRECTIONS FOR USE

BAITING: MUSKRATS, (*Ondatra zibethica*) Apply 1 to 16 CDFA Rodent Bait Blocks per bait station placement. Each CDFA Rodent Bait Block weighs approximately one ounce. Bait stations should be located near existing muskrat runways or dens. Floating bait stations may also be used. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of muskrat activity cease.

FIFRA SECTION 2(ee) RECOMMENDATION FOR USE
IN THE STATE OF CALIFORNIA

CDFA RODENT BAIT BLOCK DIPHACINONE

EPA Reg. No. 61282-26-10965

Use Directions for Wood Rats (*Neotoma* spp.)

It is a violation of federal law to use this product in any manner inconsistent with its labeling. This recommendation is permitted under FIFRA Section 2(ee). This supplemental label and the federally registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on the EPA registered label.

DIRECTIONS FOR USE

BAITING: WOOD RATS (*Neotoma* spp.) Apply 1 to 16 CDFA Rodent Bait Blocks per bait station placement. Each CDFA Rodent Bait Block weighs approximately one ounce. Bait stations should be located near existing wood rat runways or dens, Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of wood rat activity cease.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Keep away from children, pets (especially dogs), and domestic animals. Do not contaminate feed and food stuffs. Use only in rodent-infested areas where exposure to agricultural crops or commodities will not occur. Wash thoroughly with soap and water after handling and before eating or smoking. Clean clothing should be used daily.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Bait exposed to soil surfaces may be hazardous to birds and other wildlife. Do not apply directly to lakes, streams, or ponds.

STORAGE AND DISPOSAL

Store in a locked enclosure (in a properly posted area) in such a manner so as not to present a hazard to persons, animals, food, feed, crops, or property. Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Completely empty bag into application equipment. Then, dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

RODENT BAIT BLOCK DIPHACINONE TREATED GRAIN/PARAFFIN (0.005)

Active Ingredient: Diphacinone 2-diphenylacetyl-1,3-indandione 0.005%
Inert Ingredients: 99.995%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER

IMMEDIATELY: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat.

Do not induce vomiting or give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN

Intramuscular and oral administration of Vitamin K₁, combined with blood transfusions is indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin (Dicumarol).

Total: 100.000%

REGISTRANT: California Department of Food and Agriculture
1220 "N" Street, Room A357
Sacramento, CA 94271-0001

CALIF. REG. NO. 10965-50002ZA

EPA SLN. NO. CA 890021

EPA EST. NO.

NET CONTENTS: _____

DIRECTIONS FOR USE

It is a violation of State and Federal laws to use this product inconsistent with its labeling.

FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA **USE RESTRICTIONS**

For use against NORWAY RATS (*Rattus norvegicus*), ROOF RATS (*Rattus rattus*), MUSKRATS (*Ondatra zibethica*), WOOD RATS (*Neotoma* spp.) and HOUSE MICE (*Mus musculus*).

IMPORTANT

Paraffin bait blocks should only be used where environmental conditions demand a moisture-resistant bait or use of loose bait is impractical. Do not allow treated bait to contact skin.

Cereal baits imbedded in paraffin are used in sewers, outdoor placement, or other excessively damp locations where unprotected bait would spoil rapidly. Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof bait boxes. Maintain bait supply for at least 15 days. Continue baiting until all signs of feeding have stopped. Bait placements should be checked daily; replace consumed, contaminated, or spoiled baits. Where a continuous source of infestation is present, permanent bait stations should be established and the bait replenished as needed.

All poison baits must be picked up and disposed of after rodent activity has stopped. Keep this product in a closed container, free from odors which may reduce acceptability.

BAITING: NORWAY RATS, *Rattus norvegicus* and ROOF RATS, *R. rattus*.

Place a one-pound bait block in areas where rats feed, drink, or frequent (e.g., in and around barns, poultry houses, sheds, lumber and rubbish piles, and infested ditch banks).

BAITING: MUSKRATS, *Ondatra zibethica*.

Place one-pound bait blocks along ditches and waterways near muskrat burrows, runways, and other locations where fresh muskrat activity is apparent. These can be held in place by a 10-inch stake which is imbedded in the block. As blocks are consumed, replace with new ones until the desired level of control is attained.

BAITING: WOOD RATS, *Neotoma* spp. and ROOF RATS *Rattus rattus*.

Secure a bait block with a wire or nail in a tree near the damage area.

BAITING: HOUSE MICE, *Mus musculus*.

Place 4-16 ounce bait blocks in areas where mice feed, drink, or frequent (e.g., in and around barns, poultry houses, sheds, lumber and rubbish piles, and infested ditch banks).

December 94

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

Environmental Hazards

This product is toxic mammals and birds. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high-water mark.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Storage: Store only in original closed container in a cool, dry place inaccessible to children and pets. Store separately from fertilizer and away from products with strong odors which may contaminate the bait and reduce acceptability. Sweep up spillage carefully and collect for disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Commercial Use: Dispose of empty container in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **Household Use Disposal-** If empty: Do not reuse empty container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste disposal agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Read the entire Directions For Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be returned.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

Manufacturer and Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Manufacturer, and Buyer and User assume the risk of any such use. MANUFACTURER AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. THE EXCLUSIVE REMEDY OF USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES, OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF THE MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT. Manufacturer and Seller offer this product and Buyer and User accepts it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of manufacturer.



CALIFORNIA DEPARTMENT OF
FOOD AND AGRICULTURE

CDFA RODENT BAIT BLOCK DIPHACINONE

Mold and Moisture Resistant

Kills Norway rats, roof rats, and house mice in wet or dry areas

Active Ingredient:

Diphacinone [2-diphenylacetyl]-1,3-indandione].....0.005%

Inert Ingredients.....99.995%

Total:.....100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

Have label with you when obtaining treatment advice.

If Swallowed	<ul style="list-style-type: none">Call a poison control center, doctor or 1-800-498-5743 immediately for treatment advice.Have person sip a glass of water if able to swallow.Do not induce vomiting unless told to do so by the poison control center or doctor.
---------------------	---

NOTE TO PHYSICIAN

If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in bis-hydroxycoumarin overdose. Repeat as necessary based on monitoring of prothrombin times.

Distributed by: California Department of Food and Agriculture
1220 "N" Street, Room A-357
Sacramento, CA 95814

EPA Reg. No. 61282-26-10965

EPA EST. No. XXXX-CA-X

Net Contents:

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all directions and use precautions. **IMPORTANT:** Do not expose children, pets, or other non-target animals to rodenticides. To help to prevent accidents:

1. Store product not in use at a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potential destructive animals, or in areas prone to vandalism.
3. Dispose of product container. And unused, spoiled, and unconsumed baits as specified on this label.

USE RESTRICTIONS

For the control of Norway rats, roof rats, and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures. This product is especially suited for use in wet or damp areas, including river banks, gullies, irrigation ditches, sewers, garbage dumps, and landfills.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come direct contact with food. Do not apply bait directly to ground surface or in grass or other ground cover.

SELECTION OF TREATMENT AREAS

Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodent or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

Each CDFA Rodent Bait Block weighs approximately one ounce.

RATS

Apply 4 to 16 CDFA Rodent Bait Blocks (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

MICE

Place one 1-ounce CDFA Rodent Bait Block at each placement location. Space placements at 8 to 12 foot intervals. Two 1-ounce pieces may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

RATS AND MICE

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

Revised 11-18-04